



Birkeland currents in the auroral oval



Mini-groupwork 5



Current sheet 1:

$$\frac{\partial B_x}{\partial y} < 0 \Rightarrow j_z > 0$$


$$\frac{\Delta B_x}{\Delta y} \approx -\frac{15 \text{ mm}}{22 \text{ mm}} \cdot 1000 \cdot 10^{-9} = -6.8 \cdot 10^{-7} \text{ T}$$

$$\Delta y \approx \frac{10 \text{ mm}}{10 \text{ mm}} \cdot \frac{2^\circ}{360^\circ} 2\pi(R_E + 800) = 250 \cdot 10^3 \text{ m}$$

$$j_z \approx -\frac{1}{\mu_0} \frac{\Delta B_x}{\Delta y} = 2.2 \cdot 10^{-6} \text{ Am}^{-2}$$

$$j_z = -\frac{1}{\mu_0} \frac{\partial B_x}{\partial y}$$

Birkeland currents in the auroral oval




EF22445 Space Physics II

7.5 ECTS credits, P2

- shocks and boundaries in space
- solar wind interaction with magnetized and unmagnetized bodies
- sources of magnetospheric plasma
- magnetospheric and ionospheric convection
- auroral physics
- storms and substorms
- global oscillations of the magnetosphere

**First lecture Tuesday November 3, 13.15 at
Teknikringen 31, seminar room, second floor.
(Signs will be posted)**

Teknikringen 31





Thesis work at Space and Plasma Physics

Talk to Tomas



Examination

1. Written examination
(open book*), 30/10

100 p

2. Continuous examination
(mini-group works)

25 p

Grades:

A: 111-125 p

B: 96-110 p

C: 81-95 p

D: 66-80 p

E: 50-65 p

(Fx)



Written examination, 28/10, 2014, 8-13, Q21, Q26

(No academic 15 minutes!)

You may bring:

- all the course material
- any notes you have made
- pocket calculator
- mathematics and physics formula books or your favourite physics book
- formula sheet

(No computers are allowed, due to the possibility to communicate with the outside world.)

Approx. 5 different problems (which may contain sub-problems).




About the exam


Motivate your answers!

Be careful with units and
numerical calculations!

Aurora during substorm




Substorms - magnetosphere



- **GROWTH PHASE:** When IMF southward, energy is pumped into magnetotail and is stored as magnetic energy
- **ONSET:** After a certain time (~ 1 h) the magnetotail goes unstable and “snaps” due to fast reconnection.
- **EXPANSION/MAIN PHASE:** Close to Earth the magnetosphere returns to dipole-like configuration. Plasma is energized and injected into the inner parts of the magnetosphere.
- **RECOVERY PHASE:** In the outer parts of the magnetotail a *plasmoid* is ejected. The magnetosphere returns to its ground state.


Substorm Current Wedge (SCW)





Due to reconnection processes the resistivity increases here

⇒

Current takes another direction – through the ionosphere!





Substorm Current Wedge (SCW)



Auroral Electrojet (AE) index


The AE index Measures the strength of the substorm current wedge (SCW), by using the information from several magnetic observatories.



~1 – 3 h


Geomagnetic storms

Geomagnetic storms are extended periods with southward interplanetary magnetic field (IMF) and a large energy input into the magnetosphere.




Geomagnetic storms - phases

Magnetogram



Geomagnetic storms - phases




Main phase: Several particle injections increase the ring current.
Weakening of B

Initial phase: the magnetic cloud of the CME compresses the geomagnetic field.
Increase of B

Recovery phase: ring current returns to normal strength.
Recovery of B

Space weather: consequences of solar and geomagnetic activity




"conditions on the Sun and in the solar wind, magnetosphere, ionosphere and thermosphere that can influence the performance and reliability of space-borne and ground-based technological systems and can endanger human life or health."

US National Space Weather Programme

Highly energetic particles

- Particles in the radiation belts.
- Particles from solar activity (solar flares, CME)
- Cosmic radiation



Disturb or damage electronics on satellites and aeroplanes.

Danger to astronauts



Increase the rate of ionization in lower D region and thus increases absorption of radio waves.




Cosmic rays (= cosmic radiation)

Primary cosmic radiation


Extremely energetic particles ($>10^8$ eV)

- Galactic cosmic rays
- Solar 'cosmic rays' (Solar Energetic Particles)


Secondary cosmic radiation



Spectrum of galactic cosmic radiation



Simpson, 1983.



Ultra-energetic cosmic radiation.
Origin unknown. Extragalactic???

Pfotzer maximum





Fig. 1.12
Intensity profile of cosmic particles
in the atmosphere



Relativistic dynamics

Relativistic momentum

$$\mathbf{p} = \frac{m\mathbf{v}}{\sqrt{1 - \frac{v^2}{c^2}}} = \gamma m\mathbf{v}$$

$$\gamma \equiv \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

Relativistic energy

$$E = \frac{mc^2}{\sqrt{1 - \frac{v^2}{c^2}}} = \gamma mc^2$$

Relation between energy and momentum

$$E^2 = p^2 c^2 + m^2 c^4$$

Relativistic dynamics

Rest energy

$$E = mc^2$$


Kinetic energy

$$E_{kin} = E - mc^2 = mc^2 (\gamma - 1)$$

Rest energy of electron: 512 keV ~ 0.5 MeV

Rest energy of proton: 939 MeV ~ 1 GeV

!!!



24.1: Major components of the primary cosmic radiation (from Ref. 1).

Relativistic gyro radius


Non-relativistic
gyro radius

$$r_L = \frac{mv_{\perp}}{qB} = \frac{p_{\perp}}{qB}$$

Relativistic
gyro radius

$$r_L = \frac{p_{rel,\perp}}{qB} = \gamma \frac{mv_{\perp}}{qB}$$

Magnetic shielding of magnetosphere




Shielding if

$$r = \frac{p_{\perp}}{qB} < L$$


What will be the maximum energy of cosmic ray particles that will be shielded?

Interstellar plasma

Interstellar matter (10 % of Milky Way mass)



HI regions (neutral hydrogen)



HII regions
(emission nebulae)




Horsehead nebula

Triffid nebula


H1 regions are reservoirs of material for star formation

Stars are formed by gravitational collaps of interstellar clouds



Pleiades cluster


Closeup of region close to Merope



The emissions are caused by reflection by the dust particle component of the clouds.


Interstellar plasma — HII regions

- Reached by UV radiation by young hot stars.
- Mostly contains ionized hydrogen
- Approx. same density as HI regions.
- Ionization degree $\sim 100\%$
- $T \sim 10\,000\text{ K}$
- $B \sim 1\text{ nT}$



Distribution of interstellar HII gas in the Northern sky

Strömgren sphere



The size of the HII region (emission nebula) is called the Strömgren radius, R_s .

The modelled, spherical region is called a Strömgren sphere.

Strömgren radius

- The recombination rate inside the Strömgren radius is


$$r = \alpha_H n_e n_p = \alpha_H n_e^2 = \alpha_H n_H^2$$

- In equilibrium, we have

$$N_{UV} = rV = \alpha_H n_H^2 \frac{4\pi R_s^3}{3} \Rightarrow$$


$$R_s = \left(\frac{3N_{UV}}{4\pi\alpha_H n_H^2} \right)^{1/3}$$

→ Hotter star
→ Denser gas



Interstellar HI plasma

Strömgren radius



Interstellar HI plasma


N_{UV} can be determined by considering black-body radiation properties of the star (Temperature and surface area). For a hot, young star it can be $\sim 10^{49} \text{ s}^{-1}$. For a typical HI density of $n_H = 35 \text{ cm}^{-3}$, we get

$$\alpha_H \approx 3 \times 10^{-13} \text{ cm}^3 \text{ s}^{-1}$$

Red

$$R_s = \left(\frac{3N_{UV}}{4\pi\alpha_H n_H^2} \right)^{1/3} = \left(\frac{3 \cdot 10^{49}}{4\pi \cdot 3 \cdot 10^{-19} \cdot (3.5 \cdot 10^7)^2} \right)^{1/3} = 1.9 \cdot 10^{17} \text{ m} = 20 \text{ L.Y.}$$


Emission nebulae



Triffid nebula (Messier 20)



IC5146




*Heart and Soul nebulae
(IC1805, IC1848)*


- Emission nebulae often appear red, due to a prominent emission in the Balmer series
- May be non-spherical due to
 - *Gradients in the background medium*
 - *Multiple stars at the core*

Why is the chromosphere red?

Hydrogen spectrum




T_2
 T_1





H_γ H_β
434 nm 486 nm

H_α
656 nm



Interstellar magnetic field





HI regions: ~ 0.1 nT

HII regions: ~ 1 nT

Magnetic field important also in the interstellar medium!

Intergalactic matter

$2.7 \cdot 10^9$ light years



Computer simulation of intergalactic mass distribution



Intergalactic plasma

- Mostly made up of “bridges” between galaxies ($\sim 10^6$ l.y.) (Radius of Milky Way is $\sim 10^4$ l.y.)
- Detected by radio telescope measurements of synchrotron radiation from energetic electrons.
- Typical densities are 10^{-4} cm^{-3}
- Typical magnetic field: $B \sim 10^{-2} \text{ nT}$



Space research today and Space research in Sweden

The Rosetta mission to comet 67P/Churiumov-Gerasimenko




The Rosetta mission to comet 67P




3 August 2014

The Rosetta mission to comet 67P




11 September 2015


The Rosetta mission to comet 67P



Perihelion




Landing site of Philae lander




Landing Took place on November 12, 2014

Philae lander






Tail reconnection events?




Comet Encke, STEREO-A

Comets, induced magnetotail




Where does reconnection take place?

How can we measure temperatures and densities in space?



Langmuir probe



Langmuir probe
from the
Rosetta
spacecraft




Rosetta spacecraft, launched 2004. Now placed in orbit around the comet 67/P Churyumov-Gerasimenko.

Irving Langmuir



- 1881-1957
- American chemist and physicist
- Nobel prize in 1932 in chemistry
- Worked with gas discharges in vacuum tubes


Electrical potential of metal sphere in a plasma



Current balance:

$$I_e + I_i + I_{ph} \approx I_e + I_{ph} = 0$$


Maxwellian distribution of velocities in a gas



$$f(v_x) = Ce^{\frac{-mv_x^2}{2k_B T}}$$

$$C = n \sqrt{\frac{m}{2\pi k_B T}}$$

Floating potential



I positive into
the probe


$$I_e + I_{ph} = 0$$

$$I_e = -An_e e \sqrt{\frac{k_B T}{2\pi m_e}} \cdot e^{\frac{e(V - V_{pl})}{k_B T}} = -I_{ph}$$

$\Rightarrow$

$$V - V_{pl} = \frac{k_B T}{e} \ln \left(\frac{I_{ph}}{An_e e \sqrt{\frac{2\pi m_e}{k_B T}}} \right)$$

Electrical potential of metal sphere in a plasma with **bias current**



I positive into the probe


$$I_e + I_{ph} + I_b = 0$$

$$I_e = -An_e e \sqrt{\frac{k_B T_e}{2\pi m_e}} \cdot e^{\frac{e(V-V_{pl})}{k_B T_e}} = -I_{ph} - I_b$$

$\Rightarrow$


$$V - V_{pl} = \frac{k_B T_e}{e} \ln \left(\frac{I_{ph} + I_b}{An_e e \sqrt{\frac{2\pi m_e}{k_B T_e}}} \right)$$

Langmuir probe




I positive into the probe

$$V - V_{pl} = \frac{k_B T_e}{e} \ln \left(\frac{I_{ph} + I_b}{A n_e e} \sqrt{\frac{2\pi m_e}{k_B T_e}} \right)$$



Langmuir probe



I positive into
the probe

$$I_b - I_{ph} = An_e e \sqrt{\frac{k_B T_e}{2\pi m_e}} \cdot e^{\frac{e(V-V_{pl})}{k_B T_e}}$$



$\Rightarrow$

$$\ln(I_b - I_{ph}) = \ln \left(An_e e \sqrt{\frac{k_B T_e}{2\pi m_e}} \cdot e^{\frac{e(V-V_{pl})}{k_B T_e}} \right)$$

$$= \ln \left(An_e e \sqrt{\frac{k_B T_e}{2\pi m_e}} \right) + \ln \left(e^{\frac{e(V-V_{pl})}{k_B T_e}} \right) = C + \frac{e(V - V_{pl})}{k_B T_e}$$


Langmuir probe

$$\ln(I_b - I_{ph}) = C + \frac{e(V - V_{pl})}{k_B T_e}$$



The slope gives you $\frac{k_B T_e}{e}$ which is temperature in eV

Measurement of E-field in space plasmas



$$An_e e \sqrt{\frac{k_B T}{2\pi m_e}} \cdot e^{\frac{eV}{k_B T}} + I_{ph} = 0$$


Potential between plasma and probe

Since the potential between the plasma and each probe is constant if the probes are identical, the E-field can be measured by taking the potential difference between the probes and divide by the probe separation length.


$$V_3 = (V_{\text{probe},3} - V_{\text{plasma}})$$

$$(V_{\text{probe } 4} - V_{\text{probe } 3})/L = (V_4 + V_{\text{plasma } 4} - V_3 - V_{\text{plasma } 3})/L = (V_{\text{plasma } 4} - V_{\text{plasma } 3})/L = E$$

Langmuir probe and electric field probes on a spacecraft



FAST





Rosetta



Electric field measurements from Rosetta

2015-04-16









Some unsolved problems

The auroral oval is the projection of the plasmasheet onto the atmosphere



Substorms



How do they start?


- *An internal instability?
("Rubber band breaks")*

or

- *An external trigger, e.g.
IMF turns from northward
to southward?
("Rubber band released")*

Theta-aurora

How is it created?



Why is the corona so hot?





Figure 5.3. Distribution of average temperature in the solar atmosphere (Athay 1976).


Two main theories:

- 1) *Energy transported by plasma waves*
- 2) *Reconnection frees magnetic energy*

Cosmic radiation and climate




Sun spot number (R) and flux of cosmic radiation



Variations in cloud cover and cosmic radiation

Heliosphere




How is the heliosphere/heliopause created by the interaction of the solar wind and the interplanetary medium?

What is the role of the ‘termination shock’, where the solar wind is abruptly decelerated to below 100 km/s?

Swedish space research

Satellites

Rockets




Ground-based
measurements: *radar*,
optical instruments

Three IS Radars





Location

- World's only multistatic IS radar
- Three co-operating installations
- Heater
- Other instruments
 - Optical
 - Satellite
 - Radio wave




SuperDARN + ISR's






Swedish research satellites

KTH equipment in space




Viking





- Launch: 1986
- Apogee: 13500 km
- Perigee: 800km

Auroral acceleration region




Freja

The mirror image of the aurora – the black aurora




- Launched 1992
- Apogee: 1800 km
- Perigee: 600km

Freja instrumentation



E-field instrument

Solar panel




Rocket engine


Antenna

Magnetometer

Langmuir probe
(n_e measurement)





Particle instrument



UV camera


Micro satellite

Astrid II



Miniturization of the payload gave a very cheap mission


- Launched 1998
- Apogee: 1000 km
- Perigee: 1000 km



Cluster

- Launched 2000
- Apogee: $20 R_E$
- Perigee: $4 R_E$
- Separations:
200-10000 km


Future mission



Bepi-Colombo

- ESA-JAXA mission
- Launch 2013
- Arrival at Mercury: 2019
- KTH involved in MEFISTO, an electric field instrument


Future mission



Solar orbiter

- ESA mission
- Aphelion 0.28 AU (inside Mercury's orbit)
- KTH part of electric field instrument in RPW (radio and plasma waves)
- Final go-ahead 2011-10-05!
- Launch 2017-2018

Newly launched



MMS – *Magnetospheric Multiscale*

- NASA mission
- Launch 2015
- Apogee 12-31 R_E
- KTH have provide an electric field instrument – a part of the FIELDS instrument suite

My own research

Spatial or temporal variations?



t_1

Slussen




t_2

Tekniska högskolan



Interpretation 1: temporal variation

t_1



Slussen

Tekniska högskolan



t_2



Slussen

Tekniska högskolan



Interpretation 2: spatial variation

t_1



Slussen



Tekniska högskolan

t_2





Slussen




Tekniska högskolan

Interpreting Cluster multipoint measurements



Cluster field data





CLUSTER multipoint measurements (1)

E


current

plasma density




Marklund et al, 2001


Cluster passage through black aurora



Temporal evolution of acceleration potential over black aurora






Model of ionospheric evacuation by currents




Assumptions:

$$\begin{array}{l} j_{||} \text{ carried by } e^- \\ j_p \text{ carried by ions} \end{array} \quad \Rightarrow \quad \frac{\partial n_e}{\partial t} = -\frac{1}{q} \frac{\partial j_{||}}{\partial z}$$






Observations of “Enhanced Aurora”




Discrete auroral arcs sometimes exhibit intensity enhancements in thin horizontal layers at the bottom of the arc.


Enhanced Aurora theory




*Dominating process for transfer
of energy and plasma*



Plasmoids and magnetosphere



Geometry of penetrating elements unclear




A diagram showing a particle (represented by a shaded, elongated shape) moving along a curved path. The path is defined by a dashed line and a solid line. A central point is labeled 'N'. A vector $\vec{B}_e$ points upwards and to the left. A vector $\vec{B}_m$ points towards the central point 'N'. A dashed line with an arrow points to the left, labeled 'x'. A dashed line with an arrow points downwards. A shaded semi-circle is centered at 'N'.

EF2240 Space Physics 2015


Penetration mechanisms

Magnetic expulsion
(followed by localized reconnection)



Ma, 1991



Self-polarization
 $E_p = -\mathbf{v} \times \mathbf{B}$




Brenning, 2005

Estimation of scale sizes

1. Along x-direction: width of half maximum of Δn_e
2. y and z: in effect only four measurement points




2002-12-23 (39 315 s)



Method 1a



Penetration parameters



$$W_k = \frac{m_i n_e v_d^2}{2}$$

$$W_{i,th} = \frac{m_i n_e v_{i,th}^2}{2}, \quad T = 321 \text{ eV}$$

$$W_B = \frac{B_0^2}{2\mu_0}, \quad B_0 = 50 \text{ nT}$$

$$\beta_k = \frac{W_k}{W_B}$$



$$\beta_{i,th} = \frac{W_{i,th}}{W_B}$$

$$w = l_x$$


$$r_{gi} = \frac{m_i v_d}{e B_0}$$

$$K = 2.3$$


Where do they come from?



Where do they come from?



Do they exist at other planets?



MESSENGER data from Mercury

