



HOMWORK 3

Solving homework problems is an important practice and can improve your grade. It is therefore especially important that your work is:

- legible and written in an understandable way with full sentences and complete arguments,
- original, i.e. not copied or paraphrased from another source, and
- handed in on time.

Submission. The solutions can be hand-written or typed and should be submitted before the deadline, Monday November 23, 2pm. Either hand in the solutions in class, in the black mailbox for homework outside the math student office at Lindstedtsvägen 25, or by email to `boij@kth.se`. If submitted by email, the homework should be in one pdf-file and typed or scanned with high contrast.

Scoring. The maximal total score from all twelve sets of homework is 36 and the total score will be divided by nine and rounded up when counted towards the first part of the final exam. For each set of homework problems, the maximal score is 3 which corresponds to $2/3$ of the points of the problems, i.e., $\min\{3, \Sigma/2\}$.

Problem 1. Let $\Phi: G \rightarrow H$ be a group homomorphism.

- (a) Show that $\Phi(K) = \{\Phi(a) : a \in K\}$ is a subgroup of H if $K \leq G$. **(1 p)**
- (b) Show that $\Phi^{-1}(K) = \{a \in G : \Phi(a) \in K\}$ is a subgroup of G if $K \leq H$. **(1 p)**
- (c) Show that $\Phi^{-1}(K) \leq G$ is normal if K is a normal subgroup of $\Phi(G)$. **(1 p)**

Problem 2. Let $\{H_i\}_{i \in I}$ be a collection of subgroups of G such that for each pair $(i, j) \in I \times I$, we either have $H_i \subseteq H_j$ or $H_j \subseteq H_i$.

- (a) Show that $\bigcup_{i \in I} H_i$ is a subgroup of G . **(2 p)**
- (b) Show that the unions of subgroups need not be subgroups in general. **(1 p)**

Problem 3. Let $\sigma = (12)(3)(45)$ and $\tau = (135)(2)(4)$ be elements of S_5 . Determine the order of the subgroup $\langle \sigma, \tau \rangle$. **(3 p)**