



Seminar 4

See www.kth.se/social/course/SF1626 for information about how the seminars work and what you are expected to do before and during the seminars.

This seminar will start with a quiz on a variant of one of the recommended exercises from the text book *Calculus* by Adams and Essex (8th edition) which are marked by boldface in the following list:

Section	Recommended exercises
14.1	15, 19, 21
14.2	3, 5 , 15, 23
14.3	1, 3 , 13, 27
14.4	5, 9 , 15, 19 , 21
14.5	5, 7 , 9
14.6	3, 7, 11
14.7	5, 9, 13 , 21, 27

In the seminar the following problems will be discussed.

PROBLEMS

Problem 1. Let H be a regular hexagon in the xy -plane with all six vertices on the unit circle, one of which is in $(1, 0)$.

(a) Compute the integral

$$\iint_H xy \, dx dy.$$


(b) Compute the integral

$$\iint_H (x^2 + y^2) \, dx dy.$$

(c) Compute the integral

$$\iint_H (x - y)^2 \, dx dy.$$

Problem 2. A rectangular sheet of metal is shaped as the graph of the function $f(x, y) = a(\cos kx + \cos ky)$ where $a = 4,5$ mm and $k = 0,2$ mm⁻¹.



The measures of the sheet are 400 mm in the x -direction and 300 mm in the y -direction. When the sheet is placed horizontally the holes can carry some water.

- How many holes are there in the sheet?
- Use an integral in order to compute how much water the sheet can carry.

Problem 3. In order to compute an integral over a triangle in the plane it is possible to first carry out a change of variables that moves the triangle to the triangle Δ with vertices $(0, 0)$, $(1, 0)$ and $(0, 1)$. There are many variable changes that makes this possible, but the easiest way is to use an *affine* change of variables, i.e., one that is given by a linear transformation plus a constant. In other words, this can be expressed as

$$\begin{cases} x = a + bs + ct, \\ y = d + es + ft. \end{cases}$$

where a, b, c, d, e and f are constants and s and t are the new variables.

Consider the triangle T with vertices $(1, 4)$, $(2, 3)$ and $(-1, -1)$.

- Choose an affine change of variables that transforms the triangle T into the triangle Δ .
- Compute the Jacobian $\frac{\partial(x, y)}{\partial(s, t)}$ for the change of variables in (a).
- Use the change of variables in (a) in order to compute

$$\iint_T (xy - y^2) dx dy.$$