



KTH Teknikvetenskap

SF1625 Envariabelanalys
Lösningsförslag till tentamen 2016-01-11

DEL A

1. The 1st of January 2006 10 kg of a certain radioactive substance was locked up in a basement. The substance decays at a rate proportional to the amount of the substance present, with a half life of 50 years. How much of the substance remains on January 1st 2016?

Lösning. If $y(t)$ describes the amount of the substance at time t , where $t = 0$ is 1st of January 2006, then $y'(t) = ky(t)$ for some constant k and consequently $y(t) = Ce^{kt}$ where C is another constant.

At time 0 there is 10 kg of the substance, and so $C = 10$.

We know that at time $t = 50$ we have half of the original amount, that is $y(50) = 5$. We get:

$$\begin{aligned} y(50) = 5 &\iff 10e^{50k} = 5 \\ &\iff e^{50k} = \frac{1}{2} \\ &\iff 50k = \ln \frac{1}{2} \\ &\iff k = \frac{\ln \frac{1}{2}}{50} = -\frac{\ln 2}{50}. \end{aligned}$$

The amount at time t is therefore $y(t) = 10e^{-(t \ln 2)/50}$ kg. January 1st 2016 this is precisely $y(10) = 10e^{-(10 \ln 2)/50} = 10e^{-(\ln 2)/5} \approx 8.7$ kg.

□

Svar: $10e^{-(\ln 2)/5}$ kg

2. Compute the integrals:

A. $\int_0^{2\pi} |\sin x + \cos x| dx$ (hint: split the interval of integration)

B. $\int_1^e x^2 \ln x dx$ (hint: use integration by parts)

Lösning. A. Since $\sin x + \cos x$ is positive when $0 < x < 3\pi/4$, negative when $3\pi/4 < x < 7\pi/4$ and positive when $7\pi/4 < x < 2\pi$ we obtain

$$\begin{aligned} & \int_0^{2\pi} |\sin x + \cos x| dx \\ &= \int_0^{3\pi/4} (\sin x + \cos x) dx - \int_{3\pi/4}^{7\pi/4} (\sin x + \cos x) dx + \int_{7\pi/4}^{2\pi} (\sin x + \cos x) dx \\ &= 4\sqrt{2}. \end{aligned}$$

B. We use integration by parts and obtain:

$$\begin{aligned} \int_1^e x^2 \ln x dx &= \left[\frac{x^3}{3} \ln x \right]_1^e - \int_1^e \frac{x^2}{3} dx \\ &= \frac{e^3}{3} - \frac{e^3}{9} + \frac{1}{9} = \frac{2e^3 + 1}{9} \end{aligned}$$

□

Svar: $4\sqrt{2}$

3. We study the function f given by $f(x) = \frac{x+2}{x^2+1} + 2 \arctan x$.
- A. Determine the domain of definition of f .
 - B. Find the intervals where f is increasing and decreasing, respectively.
 - C. Determine whether f assumes maximum and a minimum values.
 - D. Find all asymptotes to the graph $y = f(x)$
 - E. Using the above, find the range of f .

Lösning. A. The domain is $\mathbf{R}$. Since $\lim_{x \rightarrow \pm\infty} f(x) = \pm\pi$ we have the asymptote $y = \pi$ at ∞ and the asymptote $y = -\pi$ at $-\infty$.

B. We differentiate and obtain

$$f'(x) = \frac{x^2 + 1 - 2x(x+2)}{(x^2 + 1)^2} + \frac{2}{x^2 + 1} = \frac{x^2 - 4x + 3}{(x^2 + 1)^2}.$$

We get two critical points, namely $x = 3$ and $x = 1$. We study the sign of the derivative:

If $x < 1$ then $f'(x)$ is positive.

If $1 < x < 3$ then $f'(x)$ is negative.

If $x > 3$ the $f'(x)$ is positive.

It follows that f is increasing on the interval $x \leq 1$, decreasing on the interval $1 \leq x \leq 3$ and increasing on the interval $x \geq 3$

C, D. It follows from the above that f has a local max at $x = 1$ (and $f(1) = (3 + \pi)/2$) and a local min at $x = 3$ (and $f(3) = 1/2 + 2 \arctan 3$). These local extreme values are strictly between $-\pi$ and π and hence cannot be global extreme values, because f assumes values arbitrarily close to π and $-\pi$.

To summarize, f does not assume a global max or a global min. The range is $(-\pi, \pi)$, because f is continuous and assumes values arbitrarily close to $\pm\pi$.

□

Svar: Se lösningen.

DEL B

4. We shall study Taylor approximation of the function $f(x) = \ln(1+x)$.
- A. Find the Taylor polynomial of degree 4 around the point $x = 0$ to f .
 - B. Use the polynomial from A to obtain an approximate value of $\ln 2$.
 - C. Decide whether the error of your approximation is less than 0.25.

Lösning. A. $p(x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4}$. This is standard.

B. $\ln 2 = f(1) \approx p(1) = 7/12 \approx 0.6$.

C. The error is $\frac{f^{(5)}(c)}{5!}1^5$ for some c between 0 and 1, and the absolute value of this is less than $1/5$ because $f^{(5)}(c) = 4!/(1+c)^4$. Answer yes.

□

5. We shall find the center of mass (x_T, y_T) of the upper half of the unit disc, i.e. the area given by $x^2 + y^2 \leq 1$ and $y \geq 0$. For symmetry reasons $x_T = 0$, but the y -coordinate needs to be calculated. One can show that

$$y_T = \frac{\int_0^1 2y\sqrt{1-y^2} dy}{\int_0^1 2\sqrt{1-y^2} dy}.$$

Compute the y -coordinate of the center of mass!

Lösning. The integral in the denominator gives precisely the area of the half disc, and so that integral must equal $\pi/2$ (this can also be computed, using the substitution $y = \sin x$).

The integral in the numerator can be computed thus:

$$\int_0^1 2y\sqrt{1-y^2} dy = \left[\frac{-(1-y^2)^{3/2}}{3/2} \right]_0^1 = \frac{2}{3}.$$

We see that the y -coordinate of the center of mass is $\frac{4}{3\pi}$

□

Svar: $4/3\pi$

6. An object with mass m is falling towards the surface of the earth. If we assume the air resistance to be proportional to the speed v , we obtain from Newton's second law the differential equation

$$mv'(t) = -kv(t) + mg$$

where k is a positive constant and g is the gravitational acceleration.

- A. Find the speed v at an arbitrary time t , if the object is released with zero speed at the time $t = 0$.
- B. Show that the speed cannot increase without bound and in fact will approach a limiting value after long time. Find this limiting value.

Lösning. We can write the differential equation

$$\frac{dv}{dt} + \frac{k}{m}v = g$$

The solution v has the structure $v = v_h + v_p$ where v_h is the general solution to the corresponding homogeneous equation (i.e. with right-hand-side 0) and v_p is any particular solution. We obviously can take $v_p = gm/k$. In order to find v_h we see that the characteristic equation $r + (k/m) = 0$ has solution $r = -k/m$ and hence $v_h = Ce^{-kt/m}$. Putting all this together we get:

$$v(t) = Ce^{-kt/m} + \frac{gm}{k}, \quad C \text{ an arbitrary constant,}$$

is the general solution to the differential equation. If the object is released with zero speed then $v(0) = 0$ and so we must choose $C = -gm/k$.

The speed at an arbitrary time t is therefore given by

$$v(t) = \frac{-gm}{k}e^{-kt/m} + \frac{gm}{k} = \frac{gm}{k}(1 - e^{-kt/m}).$$

B. Since $e^{-kt/m}$ is decreasing, v is increasing, and since $\lim_{t \rightarrow \infty} v(t) = gm/k$ this is the limiting value.

□

Svar: Se lösningen.

DEL C

7. This problem is about the theory of local extreme points.

- A. Define what is meant by a local maximum of a function f .
- B. Prove this statement: If the function f assumes a local maximum at an interior point a in the domain of definition and f is differentiable at a , then $f'(a) = 0$.
- C. Give an example showing that a function can have a derivative that is 0 at a point without assuming a local extreme value at that point.
- D. Give an example showing that a function can assume a local maximum at a point without having a derivative that is zero at that point.

Lösning. A. Punkten a i definitionsmängden till f är en lokal maxpunkt till f om det finns en omgivning I till a sådan att $f(a) \geq f(x)$ för alla $x \in I$ som ligger i definitionsmängden till f .

B. Anta att a i det inre av definitionsmängden är en lokal maxpunkt till f . För alla tillräckligt små positiva tal h gäller då att

$$\frac{f(a+h) - f(a)}{h}$$

är mindre än eller lika med 0, eftersom täljaren är negativ och nämnaren positiv. Det följer att $f'(a)$, som ju är gränsvärdet av detta, också måste vara mindre än eller lika med 0.

För alla tillräckligt små negativa tal h gäller tvärtom att

$$\frac{f(a+h) - f(a)}{h}$$

är större än eller lika med 0, eftersom täljaren är negativ och nämnaren negativ. Det följer att $f'(a)$, som ju är gränsvärdet av detta, också måste vara större än eller lika med 0.

Eftersom 0 är det enda tal som både är större än eller lika med 0 och mindre än eller lika med 0 så måste $f'(a) = 0$.

C. Funktionen $f(x) = x^3$ uppfyller att $f'(0) = 0$ samtidigt som $x = 0$ inte är en lokal extrempunkt.

D. Funktionen $g(x) = -|x|$ har en lokal maxpunkt i origo men har inte någon derivata där.

□

Svar: Se lösningen

8. We study the curve $y = x^4$. For each point (x, y) on the curve (except for the origin) the curve has a normal line intersecting the y -axis at exactly one point $(0, b)$. Find the smallest possible value of b .

Lösning. Put $f(x) = x^4$. Since f is even it is enough to study the problem for positive x . We differentiate and obtain $f'(x) = 4x^3$. The slope of the normal to the curve $y = x^4$ at (x_0, x_0^4) is given by $-1/4x_0^3$. The equation of the normal line is

$$y - x_0^4 = -\frac{1}{4x_0^3}(x - x_0).$$

This line cuts the y -axis at the point

$$(0, x_0^4 + \frac{1}{4x_0^2}).$$

Hence we want to minimize the function $d(x) = x^4 + \frac{1}{4x^2}$ då $x > 0$. We differentiate and get

$$d'(x) = 4x^3 - \frac{1}{2x^3}.$$

For positive x we get

$$d'(x) = 0 \iff x^6 = \frac{1}{8} \iff x = \frac{1}{\sqrt[6]{8}}$$

Observing that $d'(x)$ is negative when $0 < x < 1/\sqrt[6]{8}$ and positive when $x > 1/\sqrt[6]{8}$ we see that this point gives the minimum value.

The minimum value of b is $b = d(1/\sqrt[6]{8}) = 3/4$.

□

Svar: 3/4

9. Determine whether the improper integral

$$\int_0^{\pi} \frac{dx}{x \sin x + \sqrt{x}}$$

is convergent or divergent.

Lösning. Integral is improper at $x = 0$ because the integrand is not bounded there. We have

$$0 \leq \frac{1}{x \sin x + \sqrt{x}} \leq \frac{1}{\sqrt{x}} \text{ då } 0 < x < \pi,$$

since $x \sin x \geq 0$. Since furthermore

$$\int_0^{\pi} \frac{dx}{\sqrt{x}} = 2\sqrt{\pi}$$

it follows that

$$\int_0^{\pi} \frac{dx}{x \sin x + \sqrt{x}}$$

is convergent.

□

Svar: Convergent
