


# Minigroupwork 2, solutions, 2016


a)



TIME RANGE=2009/6/10 (161) to 2009/6/10 (161)



Please acknowledge data provider, J.H. King, N. Papatashvili at Perot Sys, NASA GSFC and CDAWeb when using these data.  
Generated by CDAWeb on Thu Sep 10 14:43:40 2009



$$\psi = \arctan \frac{\omega_{sun} r}{u_{sw}} \quad \Rightarrow \quad u_{sw} = \frac{\omega_{sun} r}{\tan \psi}$$

$$\omega_{sun} = 2\pi/T = 2.9 \cdot 10^{-6} \text{ s}^{-1} \quad (T = 25 \text{ days at equator})$$

$$r = 1 \text{ A.U.} = 1.496 \cdot 10^{11} \text{ m.}$$

$$\tan \psi = |B_y/B_x| \approx 3.6/2.6 \text{ (from figure)} \quad (\psi = 54^\circ)$$

With these figures I get  $u_{sw} = 313 \text{ km/s}$

**b)**

The magnetic Reynolds number is calculated by using typical plasma flow velocities  $v_c$  and typical length scales of magnetic field variations  $l_c$

Use solar wind velocity obtained in a) for typical flow velocity. To obtain  $l_c$ , multiply the time  $t$  it takes the magnetic field structure (indicated in the figure), to pass over the satellite and use  $l_c = vt$ . I get  $l_c = 2.8 \cdot 10^8 \text{ m}$ .

Using a temperature of  $5 \cdot 10^4 \text{ K}$ , we can evaluate the conductivity, remembering that the temperature should be given in eV. We get the conversion from

$$W = \frac{3}{2} k_B T$$

which gives the result that 1 eV corresponds to a temperature of 7729 K. We then get  $T = 6.5 \text{ eV}$ , and

$$\sigma = 3.1 \cdot 10^4 \text{ S/m}$$

Putting in the numbers I get

$$R_m = \mu_0 \sigma v_c l_c \approx 3.5 \cdot 10^{12} \gg 1$$

So the solar wind magnetic field is frozen into the plasma to a very good approximation.

**c)**

$$\rho = n_e m_p = 6.1 \cdot 10^6 \cdot 1.67 \cdot 10^{-27} = 1.02 \cdot 10^{-20}$$

Then the kinetic energy density is ( $v = 313 \text{ km/s}$ ):

$$\rho v^2 / 2 = 5 \cdot 10^{-10} \text{ Jm}^{-3}$$

The magnetic energy density is (using values of figure)

$$\frac{B^2}{2\mu_0} = \frac{B_x^2 + B_y^2 + B_z^2}{2\mu_0} = (2.6^2 + 3.6^2 + 1.7^2) \cdot (10^{-9})^2 / 2\mu_0 = 9.0 \cdot 10^{-12} \text{ Jm}^{-3}$$

The ratio between the kinetic and magnetic energy densities is approximately 50, thus the plasma motion determines the magnetic field configuration, and not the other way around.