


VEKTORANALYS

Kursvecka 3

Övningar

CYLINDRICAL COORDINATES



$P: (x_\theta, y_\theta, z_\theta)$ cartesian coord.
 $P: (\rho, \varphi, z_\theta)$ cylindrical coord.

$$\begin{cases} x = \rho \cos \varphi \\ y = \rho \sin \varphi \\ z = z \end{cases}$$

$$0 < \varphi < 2\pi$$


SURFACE ELEMENT

$$d\bar{S} = \hat{e}_\rho \rho d\varphi dz \quad (\text{on the lateral surface})$$


$$d\bar{S} = \hat{e}_z \rho d\varphi d\rho \quad (\text{on the top and bottom surfaces})$$

VOLUME ELEMENT

$$dV = \rho d\rho d\varphi dz$$



SPHERICAL COORDINATES



$P: (x_0, y_0, z_0)$ cartesian coord.

$P: (r, \theta, \phi)$ spherical coord.

$$r = \sqrt{x^2 + y^2 + z^2}$$

$$\begin{cases} x = r \cos \phi \sin \theta \\ y = r \sin \phi \sin \theta \\ z = r \cos \theta \end{cases}$$

$$0 < \phi < 2\pi$$

$$0 < \theta < \pi$$

SURFACE ELEMENT

$$d\bar{S} = \hat{e}_r r^2 \sin \theta d\theta d\phi$$

VOLUME ELEMENT

$$dV = r^2 \sin \theta dr d\theta d\phi$$

PROBLEM 1

Calculate: $\iint_S \bar{A} \cdot d\bar{S}$ where the vector field is: $\bar{A} = (x, y, z)$

and S is a cube (length 2 each side) centred in the origin.


(a) In a direct way (using the parameterization of the surface)

(b) Using the Gauss theorem

SOLUTION

$$(a) \quad \iint_S \bar{A} \cdot d\bar{S} = \sum_i \iint_{S_i} \bar{A} \cdot d\bar{S}$$

Let's start with S_1 :
1-parameterization of S_1 :



$$\left. \begin{array}{l} y = 1 \\ |x| < 1 \\ |z| < 1 \end{array} \right\} \Rightarrow \bar{r}(u, v) = (u, 1, v) \quad \begin{array}{l} u: -1 \rightarrow +1 \\ v: -1 \rightarrow +1 \end{array}$$

2- Integral calculation:

$$\int_{S_1} \bar{A} \cdot d\bar{S} = \int_u \int_v \bar{A}(\bar{r}(u, v)) \cdot \left(\frac{\partial \bar{r}}{\partial u} \times \frac{\partial \bar{r}}{\partial v} \right) du dv$$

$$\begin{cases} \frac{\partial \bar{r}}{\partial u} = (1, 0, 0) \\ \frac{\partial \bar{r}}{\partial v} = (0, 0, 1) \end{cases} \Rightarrow \left(\frac{\partial \bar{r}}{\partial u} \times \frac{\partial \bar{r}}{\partial v} \right) = (0, 1, 0)$$

$$\int_{S_1} \bar{A} \cdot d\bar{S} = \int_{-1}^1 \int_{-1}^1 (u, 1, v) \cdot (0, 1, 0) du dv = \int_{-1}^1 \int_{-1}^1 du dv = 4$$

Due to the symmetry of the problem we have: $\int_{S_i} \bar{A} \cdot d\bar{S} = 4$

$$\Rightarrow \iint_S \bar{A} \cdot d\bar{S} = \sum_i \iint_{S_i} \bar{A} \cdot d\bar{S} = 6 \cdot 4 = 24$$

(b) S is a closed surface \Rightarrow we can apply the Gauss theorem

$$\left. \begin{aligned} \iint_S \bar{A} \cdot d\bar{S} &= \iint_V \operatorname{div} \bar{A} dV \\ \operatorname{div} \bar{A} &= \frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y} + \frac{\partial A_z}{\partial z} = 1 + 1 + 1 = 3 \end{aligned} \right\} \Rightarrow \iint_S \bar{A} \cdot d\bar{S} = \iint_V 3 dV = 3V = 3 \cdot 2^3 = 24$$

PROBLEM 2


Calculate $\iint_S \bar{A} \cdot d\bar{S}$ using the Gauss theorem

where the vector field is: $\bar{A} = (x^3, y^3, z^3)$

and the surface S is a half sphere defined by:

$$\begin{cases} x^2 + y^2 + z^2 = R^2 \\ x + y \geq 0 \end{cases}$$

SOLUTION



But S is NOT a closed surface!
So we can consider the surface


$$S_{tot} = S + S_{plane}$$

$$\iint_{S_{tot}} \bar{A} \cdot d\bar{S} = \iiint_V \operatorname{div} \bar{A} dV$$
$$\iint_S \bar{A} \cdot d\bar{S} = \iint_{S_{tot}} \bar{A} \cdot d\bar{S} - \iint_{S_{plane}} \bar{A} \cdot d\bar{S}$$

$$\iint_S \bar{A} \cdot d\bar{S} = \iint_V \operatorname{div} \bar{A} dV - \iint_{S_{\text{plane}}} \bar{A} \cdot d\bar{S}$$

So we have transformed a surface integral into a volume integral minus another surface integral
What is the advantage?
They can be calculated much easier!!

Let's consider the second integral.



$$S_{\text{plane}} \quad \begin{cases} x^2 + y^2 + z^2 \leq R^2 \\ x + y = 0 \end{cases}$$

$$\text{On } S_{\text{plane}} \quad x = -y \quad \Rightarrow \quad \bar{A} = (x^3, -x^3, z^3)$$

$$\text{On } S_{\text{plane}} \quad \text{the vector is perpendicular to } \hat{n} \quad \Rightarrow \quad \iint_{S_{\text{plane}}} \bar{A} \cdot d\bar{S} = 0$$

Let's consider the first integral.

$$\iint_V \operatorname{div} \bar{A} dV \quad \text{with} \quad \operatorname{div} \bar{A} = \frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y} + \frac{\partial A_z}{\partial z} = 3x^2 + 3y^2 + 3z^2 \quad \downarrow$$

Spherical coordinates

due to symmetry

$$\text{since } \operatorname{div} \bar{A} = 3r^2 \Rightarrow \iiint_{V_{sphere}} \operatorname{div} \bar{A} dV = \frac{1}{2} \iiint_{V_{sphere}} \operatorname{div} \bar{A} dV$$

$$\iiint_{V_{sphere}} \operatorname{div} \bar{A} dV = \int_0^{2\pi} \int_0^\pi \int_0^R 3r^2 r^2 \sin \theta dr d\theta d\phi = 3 \int_0^{2\pi} d\phi \int_0^\pi \sin \theta d\theta \int_0^R r^4 dr = \frac{12\pi R^5}{5}$$

$$\Rightarrow \iiint_{V_{sphere}} \operatorname{div} \bar{A} dV = \frac{1}{2} \iiint_{V_{sphere}} \operatorname{div} \bar{A} dV = \frac{6\pi R^5}{5}$$

$$\iint_S \bar{A} \cdot d\bar{S} = \iiint_{V_{sphere}} \operatorname{div} \bar{A} dV - \iint_{S_{plane}} \bar{A} \cdot d\bar{S} = \frac{6\pi R^5}{5}$$


PROBLEM 3

Calculate the line integral of the vector field: $\bar{A} = (y + 2x, x^2 + z, y)$

along the closed curve: $\begin{cases} \bar{r}(u) = (\cos u, \sin u, f(u)) \\ u: 0 \rightarrow 2\pi \end{cases}$ with $f(0) = f(2\pi)$

- (a) directly
- (b) using the Stokes' theorem

SOLUTION



The curve is on the cylinder
defined by $(\cos u, \sin u, z)$

On the cylinder the curve
is defined by $z=f(u)$

SOLUTION (A)

We will calculate $\int_L \bar{A} \cdot d\bar{r} = \int_a^b \bar{A}(\bar{r}(u)) \cdot \frac{d\bar{r}}{du} du$

$$\frac{d\bar{r}}{du} = \left(-\sin u, \cos u, \frac{df}{du} \right)$$

$$\bar{A}(\bar{r}(u)) = (\sin u + 2 \cos u, \cos^2 u + f(u), \sin u)$$

$$\Rightarrow \int_L \bar{A} \cdot d\bar{r} = \int_0^{2\pi} \left(\sin u + 2 \cos u, \cos^2 u + f(u), \sin u \right) \cdot \left(-\sin u, \cos u, \frac{df}{du} \right) du =$$

$$= \int_0^{2\pi} \sin^2 u du - 2 \int_0^{2\pi} \sin u \cos u du + \int_0^{2\pi} \cos^3 u du + \int_0^{2\pi} \left(f(u) \cos u + \frac{df}{du} \sin u \right) du =$$

$$= \left[\frac{u}{2} - \frac{\sin 2u}{4} \right]_0^{2\pi} - \left[\sin^2 u \right]_0^{2\pi} + \left[\sin u - \frac{1}{3} \sin^3 u \right]_0^{2\pi} + \left[f(u) \sin u \right]_0^{2\pi} = -\pi$$

SOLUTION (B)

$$\int_L \bar{A} \cdot d\bar{r} = \iint_S \bar{\operatorname{rot}} \bar{A} \cdot d\bar{S}$$

$$S = S_1 + S_2$$

$$\bar{\operatorname{rot}} \bar{A} = \begin{vmatrix} \hat{e}_x & \hat{e}_y & \hat{e}_z \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ y+2x & x^2+z & y \end{vmatrix} = (1-1, 0-0, 2x-1) = (0, 0, 2x-1)$$

$\bar{\operatorname{rot}} \bar{A}$ is in the z-direction $\Rightarrow \iint_{S_1} \bar{\operatorname{rot}} \bar{A} \cdot d\bar{S} = 0$

$$\int_L \bar{A} \cdot d\bar{r} = \iint_S \bar{\operatorname{rot}} \bar{A} \cdot d\bar{S} = \iint_S (0, 0, 2x-1) \cdot \hat{e}_z dxdy = \iint_{S_2} (2x-1) dxdy$$

cylindrical coord.

$$\begin{aligned} &= \int_0^{2\pi} \int_0^1 (2\rho \cos \varphi - 1) \rho d\rho d\varphi = 2 \int_0^{2\pi} \cos \varphi d\varphi \int_0^1 \rho^2 d\rho - \int_0^1 \rho d\rho = -2\pi \left[\frac{\rho^2}{2} \right]_0^1 = -\pi \end{aligned}$$

