MATRIX ALGEBRA MAGNUS JANSSON DEADLINE: 2014–05–19, 10.00

## Homework # 7

1. Let

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$$
$$B = \begin{bmatrix} 0 & 2 \\ 0 & 0 \end{bmatrix}$$

Determine F(A), F(B), F(A+B), and show that F(A+B) is a proper subset of F(A) + F(B).

- 2. Show that  $A \in M_2(R)$  is positive stable if and only if tr(A) > 0 and det(A) > 0.
- 3. Let  $B \in M_n$  be a matrix none of whose eigenvalues  $\mu_i$  is equal to 1, and define

$$A = (B + I)(B - I)^{-1}$$

Show that the eigenvalues of A are given by

$$\lambda_i = \frac{\mu_i + 1}{\mu_i - 1}$$

Prove that  $\operatorname{Re}(\lambda_i) < 0$  if and only if  $|\mu_i| < 1$ . Conclude that A is negative stable if and only if B is a convergent matrix.

4. Use the transformation in the previous exercise to derive the discrete time version of the Lyapunov equation from the continuous time counterpart. (That is, derive  $A^*GA - G = -H$  from  $GA + A^*G = -H$ , where the last negative sign is because we discuss negative stability here.)