



Last lecture (1)

- Course info
- Definition of plasma
- Solar interior and atmosphere
- Plasma physics 1

Today's lecture (2)

- Plasma physics 2
- Solar activity



Steps to take to take the course

1) Make sure you have signed up for the course.

If you haven't: contact your Masters coordinator or studievägledare

2) Register for the course! (My Pages)

You have to do this yourself!



Examination

1. Written examination
(open book*), 30/10

100 p

2. Continuous examination
(mini-group works)

25 p

Grades:

A: 111-125 p

B: 96-110 p

C: 81-95 p

D: 66-80 p

E: 50-65 p

(Fx)



Written examination, 30/10 2014, 08.00-13.00, M33, M37, M38

You may bring:

- all the course material
- any notes you have made
- pocket calculator
- mathematics and physics formula books or your favourite physics book
- formula sheet

(No computers are allowed, due to the possibility to communicate with the outside world.)

Approx. 5 different problems (which may contain sub-problems).

The character of the problems is such that to get a high score you will have to show that you have obtained a certain course goal, e.g. to make a reasonable order of magnitude estimate or figure out a simple model for some space physics phenomenon.

Continuous examination

Mini-group works

5 mini-group works
(5×5 p = 25 p)

Approx. 1 h during Tutorials 1-5

- *A problem similar to those on the written examination is given*
- *Groups of 3 (randomized).*
- *Elect a secretary!*
- *Write down a solution!*





Litterature

- C-G. Fälthammar, "Space Physics" (compendium), 2nd Ed, Third Printing, 2001.
- Larry Lyons, "Space Plasma Physics", from *Encyclopedia of Physical Science and Technology*, 3rd edition, 2002.
- Lecture notes and extra material handed out during lectures.



Course home page

KTH Social:

<https://www.kth.se/social/course/EF2240/>

At the home page I will post new information continuously. Here you can also find lecture notes, exercises (and some solutions), etc.

Preliminary guest lecturer



Swedish astronaut Christer Fuglesang
Lecture 10

Rosetta mission

Comet 67P/Churyumov-Gerasimenko





Photo from the Giotto spacecraft, ca 1985.


Launched, Feb., 2004, arrival August 2014.
KTH involved in plasma density instrument.



The Rosetta mission to comet 67P



15 August 2014, possible landing sites for Philae lander






Today

Activity	Date	Time	Room	Subject	Litterature
L1	2/9	10-12	Q33	Course description, Introduction, The Sun 1, Plasma physics 1	CGF Ch 1, 5, (p 110-113)
L2	4/9	10-12	Q21	The Sun 2, Plasma physics 2	CGF Ch 5 (p 114-121), 6.3
L3	8/9	13-15	Q36	Solar wind, The ionosphere and atmosphere 1, Plasma physics 3	CGF Ch 6.1, 2.1-2.6, 3.1-3.2, 3.5, LL Ch III, Extra material
T1	10/9	10-12	Q33	Mini-group work 1	
L4	15/9	13-15	Q31	The ionosphere 2, Plasma physics 4	CGF Ch 3.4, 3.7, 3.8
T2	17/9	10-12	Q33	Mini-group work 2	
L5	19/9	15-17	Q31	The Earth's magnetosphere 1, Plasma physics 5	CGF 4.1-4.3, LL Ch I, II, IV.A
L6	23/9	8-10	Q31	The Earth's magnetosphere 2, Other magnetospheres	CGF Ch 4.6-4.9, LL Ch V.
T3	24/9	14-16	Q21	Mini-group work 3	
L7	29/9	11-13	Q36	Aurora, Measurement methods in space plasmas and data analysis 1	CGF Ch 4.5, 10, LL Ch VI, Extra material
T4	1/10	15-17	Q31	Mini-group work 4	
L8	2/10	15-17	Q34	Space weather and geomagnetic storms	CGF Ch 4.4, LL Ch IV.B-C, VII.A-C
L9	8/10	13-15	Q36	Interstellar and intergalactic plasma, Cosmic radiation, Swedish and international space physics research.	CGF Ch 7-9
T5	9/10	15-17	Q31	Mini-group work 5	
L10	13/10	15-17	Q33	Guest lecture (preliminary): Swedish astronaut Christer Fuglesang	
T6	16/10	10-12	Q36	Round-up	
Written examination	30/10	8-13	M33, M37, M38		

L = Lecture, T = Tutorial

Black-body radiation



Wien's displacement law

$$\lambda_{peak} = \frac{2.90 \times 10^{-3}}{T}$$

Stefan-Bolzmans law


$$J = \sigma_{SB} T^4$$

(J = total energy radiated per unit area per unit time)


Black-body good approximation for opaque bodies where emitted light is much more likely to interact with the material of the source than to escape.

The solar spectrum

Non-blackbody contributions




Sun's interior




$T = 15 \cdot 10^6 \text{ K}$
 $P = 4 \cdot 10^{26} \text{ W}$
 $(P/m \sim 1 \text{ mW/kg})$

The proton cycle



Energy transport in the sun



Transport by radiation, which interacts with the dense solar matter (scattering and absorption/re-emission).


It takes on average 200 000 years for a photon to reach the photosphere!

~1000 km


These convection cells are called *granulation*.

At the photosphere the mean free path of the photons becomes so large that they can reach directly out into space.

Transport by convection



Sun's interior



At the photosphere the mean free path of the photons becomes so large that they can reach directly out into space.

As a consequence also the temperature, and pressure drops.


$$p_{pl} = nk_B T$$

Example of exponential density variation in balance between pressure and gravity


$$\rho_m = \text{const} \cdot e^{-z/(k_B T / gm)} = \text{const} \cdot e^{-z/H}$$

Why is the chromosphere red?

Hydrogen spectrum



T₂
T₁



H_γ H_β
434 nm 486 nm


H_α
656 nm




Solar atmosphere

Chromosphere


Reddish colour due to $H\alpha$ emissions.



$H\gamma$ 434 nm $H\beta$ 486 nm $H\alpha$ 656 nm



Balmer series




Corona

White light scattered from photosphere

Total solar eclips

Hydrogen atom




$H\gamma$ 434 nm $H\beta$ 486 nm

$H\alpha$ 656 nm




Balmer series




Why is the chromosphere red?

Hydrogen spectrum




T_2
 T_1



H γ H β
434 nm 486 nm

H α
656 nm






For non-blackbody thermal light emitter (for example a thin gas) it is more complicated. Spectrum depends e.g. chemical composition, and how many atoms/molecules happen to be in state with high probability to decay and cause emission.

Energy (and wavelength) of emitted quantum can still be approximated:

Black-body radiation

$$\lambda_{peak} = \frac{2.90 \times 10^{-3}}{T}$$



Atomic energy levels


$$E \sim k_B T$$

$$E = hf$$

$$\lambda \sim \frac{hc}{k_B T}$$

Corona

- Temperature: up to 2 MK
- Density: 10^{-18} g/cm³
– 10^{-24} g/cm³
- Turns into the solar wind at high altitudes, without a sharp boundary.



The layers of the solar atmosphere





Figure 5.3. Distribution of average temperature in the solar atmosphere (Athay 1976).




Visible light ~ 6768 Å



He II emission line at 304 Å




(Fe IX/X) at 171 Å



X-ray at 0.3-5 Å

Coronal loops



What gives the loops this structure???

Coronal loops




Until next time:

Why does the plasma follow the magnetic field lines?

Corona

- Temperature: up to 2 MK
- Density: 10^{-18} g/cm³
– 10^{-24} g/cm³
- Turns into the solar wind at high altitudes, without a sharp boundary.



The layers of the solar atmosphere





Figure 5.3. Distribution of average temperature in the solar atmosphere (Athay 1976).




For non-blackbody thermal light emitter (for example a thin gas) it is more complicated. Spectrum depends e.g. chemical composition, and how many atoms/molecules happen to be in state with high probability to decay and cause emission.

Energy (and wavelength) of emitted quantum can still be approximated:

Black-body radiation

$$\lambda_{peak} = \frac{2.90 \times 10^{-3}}{T}$$



Atomic energy levels


$$E \sim k_B T$$

$$E = hf$$


$$\lambda \sim \frac{hc}{k_B T}$$




Visible light ~ 6768 Å



He II emission line at 304 Å




(Fe IX/X) at 171 Å



X-ray at 0.3-5 Å

Coronal loops



What gives the loops this structure???

Coronal loops




Why does the plasma follow the magnetic field lines?


Magnetized plasma

Extremely common in space.

In single particle description of plasma, the particles gyrate in the plane perpendicular to \mathbf{B} .




Gyro motion



Gyro motion

Consider a positively charged particle in a magnetic field.




Assume that the magnetic field is in the z-direction.

$$m \frac{d\mathbf{v}}{dt} = q\mathbf{v} \times \mathbf{B} \Rightarrow$$

$$\left\{ \begin{array}{l} m \frac{dv_x}{dt} = qv_y B \\ m \frac{dv_y}{dt} = -qv_x B \\ m \frac{dv_z}{dt} = 0 \end{array} \right. \Rightarrow \text{Constant velocity along } z$$

$$\left\{ \begin{array}{l} \frac{d^2 v_x}{dt^2} = \frac{qB}{m} \frac{dv_y}{dt} = \omega_g \frac{dv_y}{dt} = -\omega_g^2 v_x \\ \frac{d^2 v_y}{dt^2} = -\frac{qB}{m} \frac{dv_x}{dt} = -\omega_g \frac{dv_x}{dt} = -\omega_g^2 v_y \end{array} \right. \Rightarrow$$

Gyro motion



$$\begin{cases} \frac{d^2 v_x}{dt^2} = -\omega_g^2 v_x \\ \frac{d^2 v_y}{dt^2} = -\omega_g^2 v_y \end{cases}$$




$$\begin{cases} v_x = \text{Re} \left(v_{0x} e^{i(\omega_g t + \delta_x)} \right) = v_{0x} \cos(\omega_g t + \delta_x) \\ v_y = \text{Re} \left(v_{0y} e^{i(\omega_g t + \delta_y)} \right) = v_{0y} \cos(\omega_g t + \delta_y) \end{cases}$$

and

$$\begin{cases} x = \frac{v_{0x}}{\omega_g} \sin(\omega_g t + \delta_x) \\ y = \frac{v_{0y}}{\omega_g} \sin(\omega_g t + \delta_y) \end{cases}$$

Gyro motion

For a particle starting at time $t=0$ at $(x_0, 0)$ with velocity $(0, -v_\perp)$ we get (by definition $v_{0x}, v_{0y}, v_\perp > 0$).



$$\begin{cases} v_y(0) = v_{0y} \cos \delta_y = -v_\perp & \Rightarrow v_{0y} = v_\perp, \delta_y = \pi \\ v_x(0) = v_{0x} \cos \delta_x = 0 & \Rightarrow \delta_x = \pm \frac{\pi}{2} \end{cases}$$

and

$$\begin{cases} x(0) = \frac{v_{0x}}{\omega_g} \sin \delta_x = x_0 & \Rightarrow \delta_x = \frac{\pi}{2}, x_0 = \frac{v_{0x}}{\omega_g} \\ y(0) = \frac{v_\perp}{\omega_g} \sin \pi = 0 \end{cases}$$

So


$$\begin{cases} v_x = v_{0x} \cos \left(\omega_g t + \frac{\pi}{2} \right) = -v_{0x} \sin(\omega_g t) \\ v_y = v_\perp \cos(\omega_g t + \pi) = -v_\perp \cos(\omega_g t) \end{cases}$$

$$\begin{cases} x = \frac{v_{0x}}{\omega_g} \sin \left(\omega_g t + \frac{\pi}{2} \right) = \frac{v_{0x}}{\omega_g} \cos(\omega_g t) = \frac{v_{0x}}{\omega_g} \cos(-\omega_g t) \\ y = \frac{v_\perp}{\omega_g} \sin(\omega_g t + \pi) = -\frac{v_\perp}{\omega_g} \sin(\omega_g t) = \frac{v_\perp}{\omega_g} \sin(-\omega_g t) \end{cases}$$

$$\begin{cases} v_x = v_{0x} \cos(\omega_g t + \delta_x) \\ v_y = v_{0y} \cos(\omega_g t + \delta_y) \end{cases}$$

$$\begin{cases} x = \frac{v_{0x}}{\omega_g} \sin(\omega_g t + \delta_x) \\ y = \frac{v_{0y}}{\omega_g} \sin(\omega_g t + \delta_y) \end{cases}$$

Gyro motion



Then (because the force is all the time perpendicular to the velocity)

$$v_x^2 + v_y^2 = v_{0x}^2 \sin^2(\omega_g t) + v_{\perp}^2 \cos^2(\omega_g t) = v_{\perp}^2$$

so

$$v_{0x} = v_{\perp}$$

So

$$\begin{cases} x = \frac{v_{\perp}}{\omega_g} \cos(-\omega_g t) \\ y = \frac{v_{\perp}}{\omega_g} \sin(-\omega_g t) \end{cases}$$


and

$$x^2 + y^2 = \frac{v_{\perp}^2}{\omega_g^2} \equiv r_L^2 = \rho^2$$

$$\begin{cases} v_x = -v_{0x} \sin(\omega_g t) \\ v_y = -v_{\perp} \cos(\omega_g t) \end{cases}$$

$$\begin{cases} x = \frac{v_{0x}}{\omega_g} \cos(-\omega_g t) \\ y = \frac{v_{\perp}}{\omega_g} \sin(-\omega_g t) \end{cases}$$

Gyro (Larmor) radius





Magnetic force:

$$\mathbf{F} = q\mathbf{v}_{\perp} \times \mathbf{B}$$

Centripetal force:


$$\mathbf{F} = \frac{mv_{\perp}^2}{\rho} \hat{\rho}$$



$$v_{\perp} = v \cdot \sin \alpha$$

$$\rho = \frac{mv_{\perp}}{qB}$$

Gyro frequency



$$\rho = \frac{mv_{\perp}}{qB}$$


$$\omega\rho = v_{\perp}$$

\Rightarrow

$$\omega_g = \frac{qB}{m}$$

$$\omega = 2\pi f$$

Drift motion




Then

$$x = r_L \cos(-\omega_g t)$$

$$y = r_L \sin(-\omega_g t)$$

$$\omega_g = \frac{qB}{m}$$

$$r_L = \frac{mv_{\perp}}{qB}$$



Magnetized plasma

A magnetic field drastically changes some of the plasma properties because the plasma particles are tightly bound to the magnetic field lines.

It is difficult for the particles to move perpendicular to \mathbf{B} , but easy to move along \mathbf{B} .





Figure 10: Gyration of charged particle along magnetic field lines.

Think about this:



Can you think about a physical property of the plasma that varies with the direction?


(Such a property is called *anisotropic*.)

Coronal loops



Why does the plasma follow the magnetic field lines?

Gyro motion




Equipartition principle

Statistically the kinetic energy is equally distributed along the three dimensions:

$$E_{\parallel} = \frac{1}{2} k_B T$$

$$E_{\perp} = \frac{2}{2} k_B T$$

Frozen in magnetic field lines



In fluid description of plasma two plasma elements that are connected by a common magnetic field line at time t_1 will be so at any other time t_2 .

This applies if the magnetic Reynolds number is large:

$$R_m = \mu_0 \sigma l_c v_c \gg 1$$

An example of the collective behaviour of plasmas.

Maxwell's equations

Gauss' law $\nabla \cdot \mathbf{E} = \frac{\rho}{\epsilon_0}$

No magnetic monopoles $\nabla \cdot \mathbf{B} = 0$

Faraday's law $\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}$


Ampère's law $\nabla \times \mathbf{B} = \mu_0 \mathbf{j} + \mu_0 \epsilon_0 \frac{\partial \mathbf{E}}{\partial t}$

Lorentz' force equation

$$\mathbf{F} = q(\mathbf{E} + \mathbf{v} \times \mathbf{B})$$

Ohm's law

$$\mathbf{j} = \sigma \mathbf{E}$$




Energy density

$$W_B = \frac{B^2}{2\mu_0}, \quad W_E = \epsilon_0 \frac{E^2}{2}$$

$$\nabla \cdot \mathbf{A} = \frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y} + \frac{\partial A_z}{\partial z}$$

$$\nabla \times \mathbf{A} = \left(\frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z}, \frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x}, \frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} \right)$$

Field transformations (relativistic)



*Relativistic transformations
(perpendicular to the velocity u):*

$$\mathbf{E}' = \frac{\mathbf{E} + \mathbf{u} \times \mathbf{B}}{\sqrt{1 - u^2/c^2}}$$

$$\mathbf{B}' = \frac{\mathbf{B} - (\mathbf{u}/c^2) \times \mathbf{E}}{\sqrt{1 - u^2/c^2}}$$

For $u \ll c$:

$$\mathbf{E}' = \mathbf{E} + \mathbf{u} \times \mathbf{B}$$

induced
electric field

$$\mathbf{E} = \mathbf{E}' - \mathbf{u} \times \mathbf{B}$$

$$\mathbf{B}' = \mathbf{B}$$

Frozen in magnetic flux *PROOF*

$$(1) \quad \mathbf{j} = \sigma \mathbf{E}' = \sigma (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \quad \text{Ohm's law}$$

$$(2) \quad \mu_0 \mathbf{j} = \nabla \times \mathbf{B} - \frac{1}{c^2} \frac{\partial \mathbf{E}}{\partial t} \quad \text{Ampère's law}$$

$$(3) \quad \frac{\partial \mathbf{B}}{\partial t} = -\nabla \times \mathbf{E} \quad \text{Faraday's law}$$

$$(1) \Rightarrow \mathbf{E} = \frac{\mathbf{j}}{\sigma} - \mathbf{v} \times \mathbf{B}$$

$$(3+1) \Rightarrow \frac{\partial \mathbf{B}}{\partial t} = -\nabla \times \left(\frac{\mathbf{j}}{\sigma} - \mathbf{v} \times \mathbf{B} \right)$$

$$(2) \Rightarrow \frac{\partial \mathbf{B}}{\partial t} = -\nabla \times \left(\frac{\nabla \times \mathbf{B}}{\mu_0 \sigma} - \mathbf{v} \times \mathbf{B} \right)$$

$$\frac{\partial \mathbf{B}}{\partial t} = \nabla \times (\mathbf{v} \times \mathbf{B}) - \frac{1}{\mu_0 \sigma} \nabla \times (\nabla \times \mathbf{B}) =$$

$$\nabla \times (\mathbf{v} \times \mathbf{B}) - \frac{1}{\mu_0 \sigma} (\nabla (\nabla \cdot \mathbf{B}) - \nabla^2 \mathbf{B})$$

$$\therefore \frac{\partial \mathbf{B}}{\partial t} = \nabla \times (\mathbf{v} \times \mathbf{B}) + \frac{1}{\mu_0 \sigma} \nabla^2 \mathbf{B}$$

Frozen in magnetic flux *PROOF II*

$$\frac{\partial \mathbf{B}}{\partial t} = \underbrace{\nabla \times (\mathbf{v} \times \mathbf{B})}_A + \underbrace{\frac{1}{\mu_0 \sigma} \nabla^2 \mathbf{B}}_B$$

Order of magnitude estimate:

$$\frac{A}{B} = \frac{\nabla \times (\mathbf{v} \times \mathbf{B})}{\frac{1}{\mu_0 \sigma} \nabla^2 \mathbf{B}} \approx \frac{\frac{v \Delta B}{L}}{\frac{\Delta B}{\mu_0 \sigma L^2}} = v L \mu_0 \sigma \equiv R_m$$

Magnetic Reynolds number R_m :

$$R_m \gg 1 \Rightarrow \frac{\partial \mathbf{B}}{\partial t} = \nabla \times (\mathbf{v} \times \mathbf{B})$$

Frozen-in fields!


$$R_m \ll 1 \Rightarrow \frac{\partial \mathbf{B}}{\partial t} = \frac{1}{\mu_0 \sigma} \nabla^2 \mathbf{B}$$

Diffusion equation!

Frozen in magnetic flux *PROOF III*

$$R_m \gg 1 \Rightarrow \frac{\partial \mathbf{B}}{\partial t} = \nabla \times (\mathbf{v} \times \mathbf{B})$$


Consider the change of magnetic flux Φ through a surface S with contour l which follows plasma motion




$$\Phi = \int_S \mathbf{B} \cdot d\mathbf{S}$$

$$\frac{d\Phi}{dt} = \int_S \frac{\partial \mathbf{B}}{\partial t} \cdot d\mathbf{S} + \frac{d\Phi_c}{dt}$$

$\frac{d\Phi_c}{dt}$ This term is due to change in the surface S due to plasma motion




 has an area of $(\mathbf{v} \cdot dt) \times d\mathbf{l}$

The flux through  is $(\mathbf{v} \cdot dt) \times d\mathbf{l} \cdot \mathbf{B}$

$$\therefore \frac{d\Phi_c}{dt} = \int_l \mathbf{v} \times d\mathbf{l} \cdot \mathbf{B} =$$

Frozen in magnetic flux *PROOF IV*



$$\frac{d\Phi_c}{dt} = \int_l \mathbf{v} \times d\mathbf{l} \cdot \mathbf{B} =$$

$$-\int_l \mathbf{v} \times \mathbf{B} \cdot d\mathbf{l} = -\int_S \nabla \times (\mathbf{v} \times \mathbf{B}) \cdot d\mathbf{S}$$


$$\therefore \frac{d\Phi}{dt} = \int_S \frac{\partial \mathbf{B}}{\partial t} \cdot d\mathbf{S} - \int_S \nabla \times (\mathbf{v} \times \mathbf{B}) \cdot d\mathbf{S} =$$

$$\int_S \left[\frac{\partial \mathbf{B}}{\partial t} - \nabla \times (\mathbf{v} \times \mathbf{B}) \right] \cdot d\mathbf{S} = 0$$




$$\therefore \frac{d\Phi}{dt} = 0$$

Frozen in magnetic field lines



A *flux tube* is defined by following \mathbf{B} from the surface S . Due to the frozen-in theorem the flux tube keeps its identity and the plasma in a flux tube stays in it for ever.


In particular if we let the tube become infinitely thin we have the theorem of frozen-in field lines.




Frozen in magnetic field lines: some history

- Also known as *Alfvén's theorem*
- Hannes Alfvén (1908-1995), professor at KTH
- Alfvén received the Nobel prize in 1970

'for fundamental work and discoveries in magneto-hydrodynamics with fruitful applications in different parts of plasma physics'



Magnetized plasma



Solar magnetic field




Northern lights (aurora)

Different *plasma populations* (plasmas with different temperature and density) keep to their own field line, and thus “paint out” the magnetic field lines.



Coronal loop

Does the plasma follow the magnetic field (a) or the other way around (b)?



$$\beta \ll 1$$

$$\beta \gg 1$$

Depends on relative energy density (pressure)

$$p_{pl} = nk_B T$$

$$p_B = \frac{B^2}{2\mu_0}$$

$$\beta = \frac{p_{pl}}{p_B}$$

Plasma beta

$$B = 0.2 \text{ T}$$

$$n = 10^{23} \text{ m}^{-3} \text{ (~1\% of density at Earth surface)}$$

$$T = 6000 \text{ K}$$

$$\text{Plasma (thermal) pressure/energy density: } p_{pl} = nk_bT$$

$$\text{Magnetic pressure/energy density: } p_b = B^2/2\mu_0$$

$$\beta = \frac{p_{pl}}{p_B}$$



Coronal loop

Green

$\beta \gg 1$ The plasma dominates the magnetic field

Red

$\beta \sim 1$ Some complicated in-between behaviour

Blue

$\beta \ll 1$ The magnetic field dominates the plasma




Last Minute!



Last Minute!


- What was the most important thing of today's lecture? Why?
- What was the most unclear or difficult thing of today's lecture, and why?
- Other comments

Comet C/2010 X1 Elenin



- Discovered end of 2010
- Nucleus diameter: 3-4 km
- Coma diameter: 200 000 km (August, 2011)
- Long-period comet
- Perihelion: September 10, 2011, at 0.48 AU.
- Visible by eye???

Comets, Oort cloud, Kuiper belt



Comet Hale-Bopp

Comets





Figure 4.9.9. The “magnetosphere” of the comet Halley (Balsiger *et al.* 1988).

S/C Epoxi flyby of comet Hartley-2, 2011



Gyro radius





Magnetic force:

$$\mathbf{F} = q\mathbf{v}_{\perp} \times \mathbf{B}$$

Centripetal force:


$$\mathbf{F} = \frac{mv_{\perp}^2}{\rho} \hat{\rho}$$



$$v_{\perp} = v \cdot \sin \alpha$$

$$\rho = \frac{mv_{\perp}}{qB}$$

Gyro frequency



$$\rho = \frac{mv_{\perp}}{qB}$$

$$\omega\rho = v_{\perp}$$

\Rightarrow

$$\omega_g = \frac{qB}{m}$$

$$\omega = 2\pi f$$